

AD-A131 529

ON THE DISTRIBUTION OF THE STUDENTIZED MAXIMUM OF
EQUALLY CORRELATED NORM. (U) PURDUE UNIV LAFAYETTE IN
DEPT OF STATISTICS S S GUPTA ET AL. AUG 83 TR-83-31

1/1


UNCLASSIFIED

N00014-75-C-0455

F/G 12/1

NL

END
DATE
FILMED
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

ON THE DISTRIBUTION OF THE STUDENTIZED MAXIMUM
OF EQUALLY CORRELATED NORMAL RANDOM VARIABLES*


by

Shanti S. Gupta, S. Panchapakesan and Joong K. Sohn
University, Southern Illinois University and Purdue University
Technical Report #83-31


AD A 131 029

DTIC FILE COPY

PURDUE UNIVERSITY



DEPARTMENT OF STATISTICS



83 08 16 105

ON THE DISTRIBUTION OF THE STUDENTIZED MAXIMUM
OF EQUALLY CORRELATED NORMAL RANDOM VARIABLES*

by

Shanti S. Gupta, S. Panchapakesan and Joong K. Sohn
Purdue University, Southern Illinois University and Purdue University
Technical Report #83-31

Department of Statistics
Purdue University

August 1983

A

*This research was supported by the Office of Naval Research Contract
N00014-75-C-0455 at Purdue University. Reproduction in whole or in part
is permitted for any purpose of the United States Government.

ON THE DISTRIBUTION OF THE STUDENTIZED MAXIMUM OF
EQUALLY CORRELATED NORMAL RANDOM VARIABLES*

by

Shanti S. Gupta, S. Panchapakesan and Joong K. Sohn
Purdue University, Southern Illinois University and Purdue University

ABSTRACT

Let X_1, \dots, X_k have a joint k -variate normal distribution with zero means, common unknown variance σ^2 and known correlation matrix (ρ_{ij}) , where $\rho_{ij} = \rho$ for all $i \neq j$. Let s^2 be distributed independently of the X_i such that vs^2/σ^2 has a chi-squared distribution with v degrees of freedom. New tables with wider coverage and more accuracy than the previously published ones are given for the percentage points of $Y = \max_{1 \leq i \leq k} \frac{X_i}{s}$. Some basic theoretical results are given in Section 2. The next section describes Hartley's results for approximating the distribution function of Y . Besides a brief review of existing tables (Section 4), the paper discusses the construction of new tables based on Hartley's results (Section 5) and some specific applications (Section 6).

Key words and phrases: Normal variables; Equicorrelated; Studentized; Maximum; Multivariate t ; Percentage points; Tables; Subset selection procedures; Multiple comparisons; Prediction intervals; Tests of hypotheses.

AMS 1980 Classification: Primary 62H10, Secondary 62Q05.

*This research was supported by the Office of Naval Research Contract N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

ON THE DISTRIBUTION OF THE STUDENTIZED MAXIMUM OF EQUALLY CORRELATED NORMAL RANDOM VARIABLES*

by

Shanti S. Gupta, S. Panchapakesan and Joong K. Sohn
Purdue University, Southern Illinois University and Purdue University

1. INTRODUCTION

Let X_1, \dots, X_k have a joint k -variate normal distribution with zero means, a common unknown variance σ^2 and a known correlation matrix $\Omega = (\rho_{ij})$. Also, let s^2 be a random variable independent of the X_i 's such that vs^2/σ^2 has a chi-square distribution with v degrees of freedom. Then the joint distribution of t_1, t_2, \dots, t_k , where $t_i = X_i/s$, is a (central) multivariate t distribution which has the density (see Cornish [2] or Dunnett and Sobel [4])

$$(1.1) \quad g(t_1, \dots, t_k) = \frac{|A|^{\frac{1}{2}} \Gamma[(k+v)/2]}{(\sqrt{\pi})^{k/2} \Gamma(v/2)} \left[1 + \frac{1}{v} \sum_i \sum_j a_{ij} t_i t_j \right]^{-\frac{(v+k)}{2}}$$

where $A = (a_{ij}) = \Omega^{-1}$.

We are interested in the distribution of

$$(1.2) \quad Y = \max t_i = \frac{\max X_i}{s}.$$

Of course, the percentage points of the distribution of Y are the same as the corresponding equi-percentage points of the k -variate t distribution in (1.1). These percentage points are needed in several problems of statistical inference such as tests for the multiple comparisons of means (or mean vectors) against one-sided alternatives under a general analysis

*This research was supported by the Office of Naval Research Contract N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

of variance (or multivariate analysis of variance) model, simultaneous confidence intervals for multiple comparisons between several treatment means and a control mean, and selection and ranking procedures for selecting the best population and for selecting populations that are better than a control. So it is not surprising that several authors have provided tables of the percentage points of the distribution of Y . These tables differ with respect to the ranges of the various parameters, the accuracy of the values, the probability levels chosen, and the numerical methods employed in constructing these tables.

Besides providing new tables of these percentage points with wider coverage and more accuracy, our emphasis in this paper is on Hartley's method of approximating $\text{pr}\{Y \leq y\}$ by a series in which the leading term corresponds to the case of $v = \infty$ (σ^2 known) and the successive terms are in powers of v^{-1} . Hartley [11] obtained this as the solution of a certain differential-difference equation. His method is found very useful; however, no published tables of the percentage points of Y until now were based on this method.

In this paper, we discuss some basic theoretical results (Section 2) and Hartley's results with an additional correction term (Section 3). Besides reviewing briefly the existing tables (Section 4), we discuss the construction of the new tables (Section 5), and some specific applications (Section 6).

2. BASIC THEORETICAL RESULTS

In considering the distribution of Y we assume without loss of generality that $\sigma = 1$. Further, we only consider the case where $p_{ij} = p > 0$ for all $i \neq j$. In this case, it is well-known (Dunnett and Sobel [4]) that the random variables X_1, X_2, \dots, X_k can be generated from $k+1$ independent standard

normal variables Z_0, Z_1, \dots, Z_k by setting

$$(2.1) \quad X_i = (1-\rho)^{\frac{1}{2}} Z_i - \rho^{\frac{1}{2}} Z_0, \quad i = 1, \dots, k.$$

Using this representation, we can write the distribution function

$F_k(w; \rho)$ of $W = \max X_i$ in the form

$$(2.2) \quad F_k(w; \rho) = \int_{-\infty}^{\infty} \phi\left[\frac{w + \rho^{\frac{1}{2}} t}{(1-\rho)^{\frac{1}{2}}}\right] \varphi(t) dt$$

where $\varphi(\cdot)$ and $\phi(\cdot)$ denote the density and the cdf of a standard normal variate. By a similar argument, the distribution function $G_{k,v}(y; \rho)$ of Y is given by

$$(2.3) \quad G_{k,v}(y; \rho) = \int_0^{\infty} \int_{-\infty}^{\infty} \phi\left[\frac{yu + \rho^{\frac{1}{2}} t}{(1-\rho)^{\frac{1}{2}}}\right] \varphi(t) h_v(u) dt du$$

where $h_v(\cdot)$ is the density function of s . In the sequel, unless the context demands us to be more specific, we will simply write $G_v(y)$ and $F(y)$ instead of $G_{k,v}(y; \rho)$ and $F_k(y; \rho)$, respectively. It is a well-known fact that $\lim_{v \rightarrow \infty} G_v(y) = F(y)$.

The moments of Y can be written as a product of two expectations.

Let $Z_{[k]} = \max(Z_1, \dots, Z_k)$. For $r < v$,

$$(2.4) \quad \begin{aligned} E[Y^r] &= E[s^{-r}] E[(\sqrt{1-\rho} Z_{[k]} - \sqrt{\rho} Z_0)^r] \\ &= A_{-r} \sum_{j=0}^{[r/2]} \binom{r}{2j} \frac{(2j)!}{2^j j!} \rho^j (1-\rho)^{\frac{r-2j}{2}} a_{k,r-2j}, \end{aligned}$$

where $[r/2]$ is the largest integer less than or equal to $r/2$, $a_{k,i}$ is the i th moment of the largest of k independent standard normal variables, and

$$(2.5) \quad A_{\beta} = E[s^{\beta}] = \frac{\Gamma(\frac{v+\beta}{2})}{(\frac{v}{2})^{\beta/2} \Gamma(\frac{v}{2})}, \quad \beta > -v.$$

3. HARTLEY'S METHOD

Hartley [11] considers the statistic $U = W/S$, where W has a distribution involving a scale parameter σ , S is independent of W , and S^2/σ^2 has a chi-square distribution with ν degrees of freedom. In our problem, $W = \max X_i$ and $S = \sqrt{\nu} s$. We have taken $\sigma = 1$ without loss of generality. Thus $Y = \sqrt{\nu} U$. Let H_ν denote the distribution function of U . It has been shown by Hartley that

$$(3.1) \quad u \frac{d}{du} H_\nu(u) = \nu[H_{\nu+2}(u) - H_\nu(u)].$$

Hartley's method of solving the differential-difference equation (3.1) can be described briefly as follows.

Regarding the degrees of freedom ν as a continuous second variable taking on positive values, we seek a function $H(\nu, u)$ of two variables for which

$$(3.2) \quad u \frac{\partial H(\nu, u)}{\partial u} - \nu[H(\nu+2, u) - H(\nu, u)] = 0$$

and

$$(3.3) \quad \lim_{\nu \rightarrow \infty} H(\nu, u) = F(u).$$

Equation (3.2) is now converted into a partial differential equation by expanding the finite difference in a Taylor series. Taking the expansion with two more additional terms than Hartley did, we get

$$(3.4) \quad u \frac{\partial H}{\partial u} - \nu \left[2 \frac{\partial H}{\partial \nu} + 2 \frac{\partial^2 H}{\partial \nu^2} + \frac{4}{3} \frac{\partial^3 H}{\partial \nu^3} + \frac{2}{3} \frac{\partial^4 H}{\partial \nu^4} + \frac{4}{15} \frac{\partial^5 H}{\partial \nu^5} + \frac{4}{45} \frac{\partial^6 H}{\partial \nu^6} \right] = 0,$$

where the arguments of H are u and ν , and those of \bar{H} are u and some mean value between ν and $\nu+2$. By introducing as new variables $y = \log u$ and $x = \log \nu$, the partial differential equation (3.4) is transformed into

$$\begin{aligned}
(3.5) \quad \frac{\partial H}{\partial y} - 2 \frac{\partial H}{\partial x} = & 2e^{-x} \left(\frac{\partial^2 H}{\partial x^2} - \frac{\partial H}{\partial x} \right) + \frac{4}{3} e^{-2x} \left(\frac{\partial^3 H}{\partial x^3} - 3 \frac{\partial^2 H}{\partial x^2} + 2 \frac{\partial H}{\partial x} \right) + \\
& \frac{2}{3} e^{-3x} \left(\frac{\partial^4 H}{\partial x^4} - 6 \frac{\partial^3 H}{\partial x^3} + 11 \frac{\partial^2 H}{\partial x^2} - 6 \frac{\partial H}{\partial x} \right) + \\
& \frac{4}{15} e^{-4x} \left(\frac{\partial^5 H}{\partial x^5} - 10 \frac{\partial^4 H}{\partial x^4} + 35 \frac{\partial^3 H}{\partial x^3} - 50 \frac{\partial^2 H}{\partial x^2} + 24 \frac{\partial H}{\partial x} \right) + \\
& \frac{4}{45} e^{-5x} \left(\frac{\partial^6 H}{\partial x^6} - 15 \frac{\partial^5 H}{\partial x^5} + 85 \frac{\partial^4 H}{\partial x^4} - 225 \frac{\partial^3 H}{\partial x^3} + \right. \\
& \left. 274 \frac{\partial^2 H}{\partial x^2} - 120 \frac{\partial H}{\partial x} \right),
\end{aligned}$$

where the arguments of H are y and x , and those of \bar{H} are y and some mean value between x and $\log(e^x + 2)$.

Hartley obtains an approximate solution to (3.5) for large and moderate values of x with the help of an iterative process. When $x \rightarrow \infty$, the equation (3.5) tends to the limiting form $\frac{\partial H}{\partial y} - 2 \frac{\partial H}{\partial x} = 0$ to which Hartley obtains the solution $H_0(x, y) = F(e^{y+\frac{1}{2}x})$. This provides the first approximation. Expressed in terms of u and v , H_0 depends on $\sqrt{v} u$. This means that, for large v , $G_v(u) = H_v(\frac{u}{\sqrt{v}})$ is equal to $F(u)$ to a first approximation (which is independent of v). Successive closer approximations are obtained by Hartley iteratively. His method of solving the non-homogeneous partial differential equation $\frac{\partial H}{\partial y} - 2 \frac{\partial H}{\partial x} = \psi(y, x)$ uses the theory of characteristics. For details of the steps from this point on, we refer to Hartley's paper. We will just give here the results of Hartley for the three correction terms involving v^{-1} , v^{-2} , and v^{-3} , and add a fourth term involving v^{-4} which is derived by carrying out one more iteration. The details of derivation are not given here.

Let

$$(3.6) \quad G_{v,m}(y) = F(y) + \sum_{i=1}^m L_i(y)$$

where L_i is the i th correction term. Also, let $F^{(j)}(y)$ denote the j th order derivative of $F(y)$ w.r.t. y . Then the first four correction terms can be written as

$$(3.7) \quad \begin{aligned} L_1(y) &= \frac{1}{v} [\alpha^{(2)} - \alpha^{(1)}], \\ L_2(y) &= \frac{1}{6v^2} [3\alpha^{(4)} - 10\alpha^{(3)} + 9\alpha^{(2)} - 2\alpha^{(1)}], \\ L_3(y) &= \frac{1}{6v^3} [\alpha^{(6)} - 7\alpha^{(5)} + 17\alpha^{(4)} - 17\alpha^{(3)} + 6\alpha^{(2)}], \\ L_4(y) &= \frac{1}{360v^4} [15\alpha^{(8)} - 180\alpha^{(7)} + 830\alpha^{(6)} - \\ &\quad 1848\alpha^{(5)} + 2015\alpha^{(4)} - 900\alpha^{(3)} + 20\alpha^{(2)} + 48\alpha^{(1)}], \end{aligned}$$

where

$$(3.8) \quad \begin{aligned} \alpha^{(j)} &= \frac{1}{2^j} \phi^{(j)}(y), \quad j = 1, 2, \dots, 8, \\ \phi^{(1)}(y) &= y F^{(1)}(y), \\ \phi^{(2)}(y) &= y^2 F^{(2)}(y) + y F^{(1)}(y), \\ \phi^{(3)}(y) &= y^3 F^{(3)}(y) + 3y^2 F^{(2)}(y) + y F^{(1)}(y), \\ \phi^{(4)}(y) &= y^4 F^{(4)}(y) + 6y^3 F^{(3)}(y) + 7y^2 F^{(2)}(y) + y F^{(1)}(y), \\ \phi^{(5)}(y) &= y^5 F^{(5)}(y) + 10y^4 F^{(4)}(y) + 25y^3 F^{(3)}(y) + \\ &\quad 15y^2 F^{(2)}(y) + y F^{(1)}(y), \\ \phi^{(6)}(y) &= y^6 F^{(6)}(y) + 15y^5 F^{(5)}(y) + 65y^4 F^{(4)}(y) + \\ &\quad 90y^3 F^{(3)}(y) + 31y^2 F^{(2)}(y) + y F^{(1)}(y), \\ \phi^{(7)}(y) &= y^7 F^{(7)}(y) + 21y^6 F^{(6)}(y) + 140y^5 F^{(5)}(y) + \\ &\quad 350y^4 F^{(4)}(y) + 301y^3 F^{(3)}(y) + 63y^2 F^{(2)}(y) + y F^{(1)}(y), \end{aligned}$$

$$\begin{aligned}
 \phi^{(8)}(y) = & y^8 F^{(8)}(y) + 28y^7 F^{(7)}(y) + 266y^6 F^{(6)}(y) + \\
 (3.8) \quad & 1050y^5 F^{(5)}(y) + 1701y^4 F^{(4)}(y) + \\
 \text{cont.}) \quad & 966y^3 F^{(3)}(y) + 127y^2 F^{(2)}(y) + y F^{(1)}(y).
 \end{aligned}$$

The correction term $L_3(y)$ given without the details of derivation by Hartley in terms of the $\phi^{(j)}$ is found to be in error; stated in terms of the $\alpha^{(j)}$, his coefficient of $\alpha^{(3)}$ is $-22/3$ instead of -17 .

Now, it is easy to see that

$$(3.9) \quad F^{(j)}(y) = \frac{1}{j/2} \int_{-\infty}^{\infty} \phi^k \left[\frac{\sqrt{\rho}t+y}{\sqrt{1-\rho}} \right] H_j(t) \phi(t) dt,$$

where $H_j(t)$ is the Hermite polynomial in t of degree j . Thus the evaluation of $G_{v,m}(y)$ in (3.6) involves the evaluation of the integral in (3.9) for $j = 0, 1, \dots, 8$.

Before we discuss the details of our numerical evaluation of the percentage points of the distribution of Y by using the approximation $G_{v,m}(y)$ to its distribution function $G_v(y)$, we will give a brief account of the existing tables.

4. EXISTING TABLES: BRIEF ACCOUNT

As we noted earlier, several authors have discussed the distribution of Y and have provided relevant tables. We give a brief description of these tables.

Tables in the case of $v = \infty$ (i.e. the percentiles of W) have been given by Bechhofer [1], Gupta [6], Gupta, Nagel and Panchapakesan [8], and Milton [14]. Bechhofer's table gives several percentage points (ranging from 5% to 99.95%) to four decimal places for $k = 1(1)9$ and $\rho = 0.5$. Gupta [6] has tabulated the 75, 90, 95, 97.5, and 99 percentage points to three decimal places for $\rho = 0.5$ and $k = 1(1)50$. Gupta [6] has also given

the values of $F_k(w; \rho)$ to five decimal places for $k = 1(1)12$, $w = -3.5(0.1)3.5$ and seventeen selected values of ρ ranging from 0.1 to 0.9. The values of $F_k(w; \rho)$ have also been provided by Milton [14] to eight decimal places for $\rho = 0.5$, $k = 2(1)9(5)24$, and $w = 0.00(0.05)5.15$. Finally, the table of Gupta, Nagel and Panchapakesan [8] gives the 75, 90, 95, 97.5 and 99 percentage points for $k = 1(1)10(2)50$ and the seventeen values of ρ chosen by Gupta [6].

Percentage points of Y have been earlier tabulated by several authors. Pillai and Ramachandran [15] have tabulated the 95 percent points to two decimal places for $\rho = 0$, $k = 1(1)8$, and $v = 3(1)10, 12, 14, 15, 16, 18, 20, 24, 30, 40, 60, 120$ and ∞ .

Dunnett and Sobel [4] have considered only the bivariate t ($k=2$) and tabulated the percentage points (three decimal places) as well as the values of $G_{k,v}(y)$ (five decimal places) for $\rho = \pm 0.5$ and $v = 1(1)30(3)60(15)120, 150, 300, 600, \infty$. Their table of the values of $G_{k,v}(y)$ covers $y = 0.00(0.25)2.50(0.50)10.00$ and the percentage levels chosen for the other table are 50, 75, 90, 95 and 99.

Dunnett [3] and Gupta and Sobel [9] have considered the case of $\rho = 0.5$ and $k \geq 2$. Dunnett's table gives the 95 and 99 percent points to two decimal places for $k = 1(1)9$ and $v = 5(5)25$ whereas in the Gupta-Sobel table (correct to two decimal places) $k = 2, 5, 10(1)16, 18, 20(5)40, 50$; $v = 15(1)20, 24, 30, 36, 40, 48, 60, 80, 100, 120, 360, \infty$ and the percentage levels are 75, 90, 95, 97.5, 99, 99.75, 99.9, and 99.95. Dunnett's method is to numerically evaluate $G_{k,v}(y)$ for three successive values of y differing by 0.1 such that the desired value of the probability level is bracketed and then determine the percentage point by inverse interpolation.

The Gupta-Sobel table is for the percentage points of $\sqrt{2} Y$ and they use the Cornish-Fisher expansion with four adjustment terms.

Krishnaiah and Armitage [13] have tabulated the 95 and 99 percent points for $k = 1(1)10$, $v = 5(1)35$, and $\rho = 0(0.1)0.9$. They first evaluated $G_{k,v}(y; \rho)$ in (2.3) for $y = 0.1(0.20)6.1$ for the different values of ρ , v , and k using the 40-point Gauss-Hermite quadrature formula for the inner integral and 48-point Gauss quadrature formula for the outer integral. The required percentage points were then obtained by using cubic interpolation.

5. CONSTRUCTION OF THE NEW TABLE

The upper 100α percent point of the distribution of Y is denoted by $y_\alpha^* \equiv y^*(\alpha, k, v, \rho)$. Its numerical value is obtained as the solution of

$$(5.1) \quad G_{v,m}(y) = 1 - \alpha.$$

The secant method was used to find a value of y such that

$$(5.2) \quad |G_{v,m}(y) - 1 + \alpha| \leq \epsilon$$

with $\epsilon = 10^{-8}$. This value of y satisfying (5.2) is our solution y_α^* .

Now, the numerical evaluation of $G_{v,m}(y)$ involves the evaluation of the integrals $F^{(j)}(y)$ given by (3.9) for $j = 0, 1, \dots, 8$. This was accomplished by using one of two different methods, labeled Method 1 and Method 2, depending on the value of ρ ; the only exception is when $v = \infty$ in which case Method 2 was used for all the selected values of ρ . Method 1 evaluates $F^{(j)}(y)$ using the 60-point Gauss-Hermite quadrature formula. We will denote this numerically evaluated quantity by $\tilde{F}^{(j)}(y)$. To describe Method 2, letting

$$(5.3) \quad A = \rho^{\frac{1}{2}}(1-\rho)^{-\frac{1}{2}}, \quad B = (1-\rho)^{-\frac{1}{2}},$$

and changing the variable by the transformation $u = At + By$, we can write (3.9) as

$$(5.4) \quad F^{(j)}(y) = \int_{-\infty}^{\infty} \phi^k(u) H_j\left(\frac{u-By}{A}\right) \varphi\left(\frac{u-By}{A}\right) \left(\frac{B}{A}\right)^j \frac{1}{A} du.$$

The value of this integral is approximated by

$$(5.5) \quad \tilde{F}^{(j)}(y) = \int_{-9}^{9A+4B} \phi^k(u) H_j\left(\frac{u-By}{A}\right) \varphi\left(\frac{u-By}{A}\right) \left(\frac{B}{A}\right)^j \frac{1}{A} du.$$

and the integration was carried out by Gauss's method over intervals of length $D = 0.5$ starting from -9 until $9A + 4B$ was included. Note that, for convenience, the approximation for $F^{(j)}(y)$ is denoted by $\tilde{F}^{(j)}(y)$ for either method. Whatever the method, $\tilde{F}^{(j)}(y)$ is evaluated as a sum of the form $\sum_{\ell} F_{\ell}^{(j)}(t_{\ell}; y)$. While evaluating the terms of this sum, we used the IMSL subroutine for the values of the standard normal cdf. If the exponent in $-t_{\ell}^2/2$ was less than -50 [i.e., $|t_{\ell}| > 10$], the value of $F_{\ell}^{(j)}(t_{\ell}; y)$ was taken to be zero because for the ranges of ρ and v in our table, the value of the term was not more than 10^{-10} . Also, if the value of the standard normal cdf involved in the expression was less than 10^{-5} , then for $k \geq 8$, the value of $F_{\ell}^{(j)}(t_{\ell}; y)$ was taken to be zero since the value was not more than 10^{-20} for all the ρ values of the table.

Now, Method 2 is more precise than Method 1; however, it is also more expensive, the cost ratio being 6 to 1 in some extreme cases. In order to decide on the method and the number of correction terms for the range of selected values of ρ and v , some preliminary comparative studies were made.

In the case of $k = 19$ and $\alpha = 0.01$, we computed the percentage point y^* using both methods with $m = 4$ correction terms for $v = 15$ and 120 . The results were as follows.

Table 1. Upper 1% point of the distribution of Y

 $k = 19, m = 4$

ρ	v	Method 1	Method 2
0.2	15	3.96044429	3.96044429
	120	3.34250119	3.34250119
0.3	15	3.90335614	3.90335614
	20	3.32690446	3.32690446
0.5	15	3.78170980	3.78170981
	120	3.26561715	3.26561715
0.6	15	3.70147425	3.70147458
	120	3.21375097	3.21375094
0.7	15	3.59638560	3.61431834
	120	3.14096412	3.14096740

Based on this comparison, Method 1 was used for $\rho = 0.2(0.1)0.6$, and Method 2 for $\rho = 0.7(0.1)0.9$. In the same case ($k = 19, \alpha = 0.01$), the following table gives the percentage point y^* for $\rho = 0.1$ using both methods with 4 correction terms.

Table 2. Upper 1% point of the distribution of Y

 $k = 19, m = 4, \rho = 0.1$

v	Method 1	Method 2
15	4.01116552	4.01119416
30	3.60648913	3.60658967
48	3.47577569	3.47577575
60	3.43358259	3.43358261
120	3.35129394	3.35129394

Based on this, it was decided to use, in the case of $\rho = 0.1$, Method 2 for $v \leq 24$ and Method 1 for $v \geq 30$.

To illustrate the difference between using three and four correction terms, we give below the upper 1 percent point in the case of $k = 17$ and $\rho = 0.7$ for $v = 48, 60$, and 120 using Method 2.

Table 3. Upper 1% point of the distribution of Y

$$k = 17, \rho = 0.7$$

ν	3 terms	4 terms
48	2.94535892	2.94537526
60	2.92086206	2.92086933
120	2.87288348	2.87288399

For a high value of ρ , three terms seem to be quite adequate for $\nu \geq 60$.

For small values of ρ , we used three terms for $\nu = 120$ and four terms for other selected values of ρ which are all less than or equal to 60.

We summarize below the ranges of ρ , k , ν , and α for which the percentage points of Y are given in Table 4 along with information regarding the method (i.e., Method 1 or 2) and the number of correction terms (m) used. In Table 4, it should be noted that $P^* = 1 - \alpha$. The entries in the table are accurate to five decimal places.

The coverage of Table 4

(i) $\rho = 0.1$; $k = 1(1)9(2)19$; $\alpha = 1 - P^* = 0.01, 0.05, 0.10, 0.25$

ν	Method	m
15(1)20,24	2	4
30,36,48,60	1	4
120, ∞	1	3

(ii) $\rho = 0.2(0.1)0.6$; $k = 1(1)9(2)19$; $\alpha = 1 - P^* = 0.01, 0.05, 0.10, 0.25$; Method 1

ν	m
15(1)20,24,30,36,48,60	4
120, ∞	3

(iii) $\rho = 0.7(0.1)0.9$; $k = 1(1)9(2)15$; $\alpha = 1 - P^* = 0.05, 0.10$; Method 2

ν	m
15,17,20,24,36	4
60,120, ∞	3

Table 4. Upper 100 α percentage point of the studentized maximum of
equicorrelated standard normal variables.

v	P^*	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
15.	.75	.69120	1.12019	1.26136	1.51707	1.63337	1.72544	1.80152	1.86617	1.92229	2.01602	2.09234	2.15557	2.21192	2.26052
	.90	1.30661	1.72938	1.92410	2.07805	2.18708	2.27461	2.34758	2.41017	2.46734	2.55847	2.63183	2.69558	2.75072	2.79919
	.95	1.73305	2.11394	2.31577	2.45455	2.56035	2.64519	2.71801	2.77923	2.83227	2.92260	2.99503	3.05784	3.11102	3.15753
	.99	2.60240	2.93763	3.13605	3.28497	3.36910	3.45409	3.52607	3.58667	3.64421	3.73932	3.82077	3.89138	3.95429	4.01119
15.	.75	.69103	1.12031	1.26135	1.51743	1.63383	1.72041	1.79595	1.85933	1.91610	2.00920	2.08504	2.14891	2.20375	2.25195
	.90	1.30678	1.71741	1.92584	2.08009	2.17627	2.25282	2.31933	2.37566	2.43085	2.54120	2.61558	2.67851	2.73296	2.78087
	.95	1.73336	2.10937	2.31169	2.45169	2.55838	2.64393	2.71694	2.77823	2.83121	2.92155	2.99373	3.05605	3.10970	3.15423
	.99	2.59344	2.91612	3.11427	3.25393	3.33816	3.41304	3.48804	3.54874	3.60240	3.69434	3.77164	3.83664	3.89798	3.95140
17.	.75	.69320	1.12353	1.26455	1.51800	1.63433	1.72089	1.79643	1.85981	1.91658	2.00967	2.08550	2.14937	2.20421	2.25241
	.90	1.30933	1.71996	1.92839	2.08264	2.17882	2.25537	2.32188	2.37821	2.43339	2.54374	2.61812	2.68105	2.73590	2.78381
	.95	1.73581	2.09310	2.29543	2.43543	2.54213	2.62768	2.69419	2.75548	2.81155	2.90189	2.97323	3.03555	3.08970	3.13423
	.99	2.56630	2.89055	3.08889	3.22855	3.31278	3.38833	3.46419	3.53046	3.58713	3.69747	3.77011	3.83445	3.89119	3.94020
18.	.75	.68836	1.12104	1.26206	1.50930	1.62133	1.71207	1.78700	1.85064	1.90534	1.99758	2.07255	2.13558	2.19025	2.23784
	.90	1.30039	1.70634	1.91221	2.06246	2.15816	2.23340	2.31412	2.37461	2.42738	2.51611	2.58687	2.65042	2.70368	2.75055
	.95	1.73497	2.08473	2.28537	2.41117	2.51651	2.59330	2.66772	2.72871	2.77827	2.86505	2.93625	2.99845	3.04847	3.09418
	.99	2.55256	2.87170	3.05530	3.18140	3.27941	3.35900	3.42603	3.48335	3.53455	3.62135	3.69433	3.75663	3.81138	3.86031
19.	.75	.68792	1.12244	1.26346	1.50410	1.61613	1.70333	1.78320	1.84557	1.90154	1.99327	2.06729	2.13061	2.18461	2.23195
	.90	1.32775	1.70233	1.90632	2.04531	2.15103	2.23531	2.30546	2.36543	2.41775	2.50567	2.57776	2.63872	2.69147	2.73791
	.95	1.72913	2.07727	2.27051	2.40373	2.50303	2.58565	2.65465	2.71302	2.75404	2.84553	2.92042	2.98007	3.03165	3.07701
	.99	2.55547	2.85453	3.03422	3.15573	3.25644	3.33451	3.40036	3.45720	3.50722	3.59227	3.65303	3.72372	3.77691	3.82434
20.	.75	.68335	1.12059	1.26161	1.50171	1.61373	1.70093	1.77980	1.84293	1.89798	1.98904	2.06334	2.12579	2.17954	2.22666
	.90	1.32524	1.69640	1.90143	2.03987	2.14439	2.22803	2.29770	2.35722	2.40913	2.48534	2.55722	2.62827	2.68056	2.72659
	.95	1.72472	2.07360	2.26522	2.39453	2.49377	2.57553	2.64256	2.70077	2.75128	2.83635	2.90519	2.96530	3.01645	3.06146
	.99	2.52727	2.82340	3.00165	3.11497	3.20554	3.28233	3.34745	3.40337	3.44823	3.56602	3.63536	3.69472	3.74666	3.79287
24.	.75	.68405	1.11945	1.26047	1.49958	1.60947	1.69357	1.76907	1.83144	1.88551	1.97570	2.04900	2.11059	2.16358	2.21001
	.90	1.31784	1.67916	1.88345	2.02122	2.12516	2.20450	2.27342	2.33152	2.38215	2.46715	2.53675	2.59558	2.64642	2.69117
	.95	1.71068	2.04976	2.24372	2.37520	2.47269	2.54103	2.60640	2.66239	2.71131	2.79368	2.85132	2.91861	2.96922	3.01191
	.99	2.49216	2.79159	2.95167	3.06028	3.17102	3.24448	3.30614	3.35925	3.40586	3.48482	3.55015	3.60566	3.65442	3.69746

The entry in each case is the value of y^* satisfying $G_{k,v}(y^*, \rho) = P^* = 1 - \alpha$, where

$G_{k,v}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

 $\rho = .1$ (cont.)

v	P^*	K	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.	
30.	.75		.62276	1.11194	1.32655	1.48532	1.59765	1.63578	1.75842	1.82904	1.87344	1.95245	2.03476	2.05548	2.14771	2.19346
	.90		1.31042	1.67334	1.81321	2.00255	2.10305	2.18303	2.24953	2.30624	2.35532	2.43945	2.50621	2.56343	2.61287	2.65636
	.95		1.65723	2.02322	2.21152	2.33650	2.43126	2.50722	2.57053	2.62473	2.67215	2.75167	2.81702	2.87235	2.92026	2.96247
	.99		2.45722	2.74566	2.90653	3.02167	3.10817	3.17309	3.23670	3.28712	3.33133	3.40609	3.46763	3.52036	3.56606	3.60649
35.	.75		.63137	1.10895	1.32240	1.48125	1.59183	1.67930	1.75137	1.81249	1.86544	1.95366	2.02531	2.08546	2.13718	2.18247
	.90		1.30551	1.66541	1.83533	1.99076	2.06955	2.15333	2.23380	2.28553	2.33316	2.41557	2.48612	2.54228	2.59080	2.63345
	.95		1.65620	2.01576	2.15515	2.31784	2.41064	2.45505	2.54701	2.60002	2.64623	2.72407	2.78768	2.84188	2.88863	2.92981
	.99		2.45445	2.71582	2.87551	2.95353	3.06726	3.13486	3.19149	3.24016	3.28281	3.35458	3.41432	3.46486	3.50873	3.54762
40.	.75		.67334	1.10524	1.32725	1.47425	1.58450	1.67125	1.74262	1.80311	1.85549	1.94273	2.01355	2.07297	2.12405	2.16877
	.90		1.33344	1.65324	1.84651	1.97551	2.07266	2.15022	2.21436	2.26502	2.31657	2.39521	2.46127	2.51613	2.56349	2.60511
	.95		1.67722	1.99516	2.17455	2.35473	2.36526	2.43775	2.51805	2.56560	2.61455	2.69007	2.75155	2.80425	2.84957	2.88944
	.99		2.46552	2.67505	2.83143	2.95351	3.01726	3.08205	3.13625	3.18279	3.22553	3.29230	3.34856	3.39576	3.43337	3.47578
50.	.75		.67260	1.10300	1.32417	1.47120	1.58027	1.66545	1.73739	1.79750	1.84555	1.93520	2.00651	2.06550	2.11619	2.16057
	.90		1.22522	1.64535	1.85532	1.96632	2.05294	2.13443	2.20280	2.25873	2.30373	2.38233	2.44649	2.50058	2.54725	2.58825
	.95		1.67065	1.98931	2.10333	2.22105	2.37023	2.44153	2.50050	2.55158	2.59575	2.68292	2.73067	2.78201	2.82642	2.86549
	.99		2.55012	2.63740	2.80574	2.90951	2.93789	3.03102	3.10380	3.14908	3.18871	3.25555	3.31057	3.35727	3.39780	3.43358
120.	.75		.67654	1.09863	1.31305	1.46374	1.57168	1.65890	1.72700	1.78626	1.83772	1.92320	1.99250	2.05061	2.10052	2.14419
	.90		1.29223	1.63750	1.82220	1.94924	2.04332	2.11807	2.17954	2.23259	2.27833	2.35463	2.41721	2.46975	2.51505	2.55481
	.95		1.65573	1.96511	2.10512	2.25473	2.34061	2.40973	2.46711	2.51603	2.55872	2.63022	2.68870	2.73808	2.78074	2.81825
	.99		2.55782	2.61505	2.75771	2.85557	2.93052	2.93045	3.04046	3.08332	3.12777	3.16355	3.23569	3.27962	3.31771	3.35129
∞	.75		.67449	1.08423	1.31211	1.45834	1.55317	1.64742	1.71668	1.77525	1.82558	1.91027	1.97855	2.03577	2.08489	2.12785
	.90		1.28155	1.62581	1.80830	1.93201	2.02307	2.09702	2.15739	2.20871	2.25526	2.32769	2.38830	2.43928	2.48320	2.52173
	.95		1.64455	1.95622	2.11515	2.22782	2.31157	2.37849	2.43355	2.48125	2.52240	2.59126	2.64733	2.69456	2.73589	2.77185
	.99		2.52655	2.57553	2.71043	2.80409	2.87497	2.93183	2.97950	3.01972	3.05509	3.11457	3.16355	3.20464	3.24037	3.27185

The entry in each case is the value of y^* satisfying $G_{k,v}(y^*, \rho) = p^* = 1 - \alpha$, where

$G_{k,v}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

 $\rho = .2$

v	P^*	K	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
15.	.75		.63120	1.11009	1.32553	1.48440	1.59531	1.68320	1.75571	1.81729	1.87061	1.95957	2.03185	2.09255	2.14475	2.19048
	.90		1.24031	1.71349	1.81737	2.05884	2.16185	2.23615	2.31629	2.37826	2.42957	2.51649	2.58856	2.64553	2.70239	2.74890
	.95		1.75595	2.10522	2.30248	2.43320	2.51148	2.62458	2.69396	2.75335	2.89528	2.89233	2.96398	3.02428	3.07628	3.12189
	.99		2.60210	2.93343	3.12243	3.25444	3.33577	3.43793	3.50719	3.55762	3.61976	3.70561	3.74527	3.85054	3.90835	3.96044
16.	.75		.69013	1.11013	1.32248	1.48071	1.59110	1.67858	1.75070	1.81152	1.86498	1.95345	2.02532	2.08567	2.13757	2.18302
	.90		1.32876	1.72713	1.86959	2.04728	2.15158	2.23493	2.30437	2.36366	2.41536	2.50230	2.57337	2.63355	2.68563	2.73148
	.95		1.74569	2.05511	2.25949	2.42338	2.53525	2.60719	2.67559	2.73415	2.78537	2.87146	2.94203	3.00165	3.05316	3.09838
	.99		2.58244	2.90364	3.09407	3.22355	3.32286	3.40335	3.47099	3.52935	3.58068	3.65600	3.74074	3.80327	3.85828	3.90752
17.	.75		.65520	1.10890	1.32550	1.47747	1.58741	1.67449	1.74630	1.80724	1.86004	1.94807	2.01958	2.07982	2.13124	2.17645
	.90		1.32333	1.76185	1.86230	2.03923	2.14257	2.22522	2.29394	2.35255	2.40382	2.48974	2.56011	2.61960	2.67105	2.71634
	.95		1.72851	2.06574	2.27258	2.41042	2.51106	2.59300	2.65953	2.71759	2.76793	2.85255	2.92259	2.98165	3.03262	3.07743
	.99		2.58630	2.93604	3.08524	3.15648	3.23403	3.37302	3.43935	3.49551	3.54672	3.63193	3.70266	3.76323	3.81629	3.86356
18.	.75		.68216	1.10717	1.32742	1.47460	1.58414	1.67088	1.74240	1.80308	1.85568	1.94331	2.01449	2.07425	2.12564	2.17063
	.90		1.33059	1.85375	1.85823	2.03213	2.13463	2.21631	2.28475	2.34294	2.39365	2.47877	2.54846	2.60734	2.65825	2.70305
	.95		1.72407	2.07747	2.26792	2.39901	2.49857	2.57850	2.64537	2.70257	2.75253	2.83638	2.90554	2.96387	3.01432	3.05870
	.99		2.55223	2.85734	3.04735	3.17259	3.26856	3.34625	3.41144	3.46758	3.51665	3.60055	3.65951	3.72556	3.78017	3.82602
19.	.75		.62732	1.10582	1.32530	1.47204	1.58122	1.66766	1.73893	1.79938	1.85176	1.93906	2.00955	2.06955	2.12063	2.16543
	.90		1.32773	1.82241	1.83073	2.02576	2.12756	2.20895	2.27658	2.33422	2.38462	2.46904	2.53812	2.59647	2.64591	2.69128
	.95		1.72513	2.07013	2.25888	2.38868	2.48747	2.56671	2.63280	2.68940	2.73984	2.82202	2.89026	2.94800	2.99796	3.04192
	.99		2.53347	2.85090	3.02791	3.15136	3.24592	3.32344	3.38663	3.44188	3.49035	3.57240	3.64027	3.69814	3.74860	3.79335
20.	.75		.68855	1.10423	1.32340	1.46974	1.57860	1.66477	1.73581	1.79606	1.84825	1.93524	2.00507	2.06516	2.11613	2.16076
	.90		1.32534	1.82851	1.83581	2.02006	2.12123	2.20209	2.26926	2.32650	2.37555	2.46033	2.52858	2.58676	2.63679	2.68079
	.95		1.72472	2.06556	2.25059	2.37982	2.47756	2.55608	2.62156	2.67763	2.72550	2.80898	2.87657	2.93377	2.98326	3.02683
	.99		2.52797	2.83578	3.01055	3.13238	3.22567	3.30114	3.36444	3.41680	3.46666	3.54747	3.61423	3.67111	3.72063	3.76451
24.	.75		.68485	1.09934	1.31741	1.46251	1.57036	1.65568	1.72598	1.78553	1.83721	1.92321	1.99302	2.05160	2.10195	2.14602
	.90		1.31784	1.87327	1.87337	2.02119	2.10140	2.18061	2.24635	2.30244	2.35125	2.43311	2.50001	2.55646	2.60521	2.64807
	.95		1.71032	2.04301	2.22601	2.35151	2.44658	2.52387	2.58644	2.64084	2.68322	2.76818	2.83367	2.89009	2.93705	2.97927
	.99		2.45216	2.79872	2.95246	3.07316	3.16244	3.23459	3.29507	3.34706	3.39263	3.46963	3.53316	3.58716	3.63410	3.67558

The entry in each case is the value of y^* satisfying $G_{k,v}(y^*, \rho) = P^* = 1 - \alpha$, where

$G_{k,v}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

		$\rho = .2$ (cont.)																
		1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.			
30.	.75	.68276	1.09550	1.31147	1.45535	1.56220	1.64637	1.71624	1.77521	1.82525	1.91128	1.98025	2.03812	2.08785	2.13137			
	.90	1.31042	1.68419	1.85915	1.98450	2.08186	2.15945	2.22350	2.27866	2.32539	2.40334	2.47164	2.52571	2.57424	2.61800			
	.95	1.69725	2.02202	2.21190	2.32376	2.41622	2.49033	2.55203	2.60479	2.65081	2.72216	2.79155	2.84516	2.89155	2.93238			
	.99	2.45725	2.74255	2.95579	3.01544	3.10072	3.18953	3.27224	3.27679	3.32018	3.39347	3.45388	3.50519	3.54976	3.58911			
36.	.75	.68137	1.09232	1.30754	1.45061	1.55679	1.64071	1.70980	1.76834	1.81801	1.90337	1.97180	2.02919	2.07850	2.12165			
	.90	1.30551	1.68220	1.85112	1.97293	2.06900	2.14552	2.20896	2.26301	2.31002	2.38872	2.45258	2.50714	2.55387	2.59491			
	.95	1.68830	2.00957	2.18542	2.30555	2.39631	2.46900	2.52947	2.58115	2.62622	2.70191	2.76392	2.81634	2.86167	2.90156			
	.99	2.43245	2.71310	2.89949	2.97785	3.06051	3.12720	3.18300	3.23053	3.27268	3.34371	3.40205	3.45159	3.49460	3.53257			
48.	.75	.67984	1.09304	1.30722	1.44472	1.55008	1.63331	1.70179	1.75930	1.81000	1.89355	1.96129	2.01808	2.06687	2.10955			
	.90	1.29944	1.64622	1.82272	1.95065	2.05309	2.12330	2.19060	2.24365	2.28377	2.36694	2.42990	2.48294	2.52868	2.56884			
	.95	1.67722	1.99222	2.15560	2.23313	2.37179	2.44273	2.50169	2.55205	2.59593	2.66559	2.72988	2.78081	2.82484	2.86357			
	.99	2.40658	2.67620	2.82755	2.93188	3.01134	3.07536	3.12888	3.17481	3.21499	3.28277	3.33855	3.38589	3.42696	3.46320			
60.	.75	.67820	1.08630	1.29975	1.44121	1.54608	1.62829	1.69702	1.75471	1.80463	1.88768	1.95501	2.01145	2.05992	2.10232			
	.90	1.28582	1.64053	1.82535	1.95013	2.04354	2.11807	2.17959	2.23215	2.27774	2.35333	2.41618	2.46855	2.51370	2.55334			
	.95	1.67065	1.98552	2.15566	2.23984	2.35727	2.42717	2.48583	2.53481	2.57799	2.65044	2.70972	2.75977	2.80302	2.84106			
	.99	2.55012	2.65511	2.80237	2.90485	2.98243	3.04439	3.09707	3.14182	3.18055	3.24503	3.30119	3.34722	3.38713	3.42235			
120.	.75	.67654	1.08285	1.29597	1.43423	1.53813	1.62012	1.68753	1.74459	1.79355	1.87602	1.94253	1.99625	2.04609	2.08793			
	.90	1.28385	1.62830	1.81077	1.93227	2.02494	2.09782	2.15811	2.20939	2.25353	2.32536	2.38501	2.44005	2.48403	2.52261			
	.95	1.66765	1.98435	2.15470	2.24365	2.32864	2.39349	2.45280	2.50632	2.55463	2.61253	2.66505	2.71286	2.75937	2.79563			
	.99	2.55722	2.61095	2.75462	2.85201	2.92594	2.98535	3.03491	3.07756	3.11446	3.17691	3.22820	3.27166	3.30931	3.34250			
∞	.75	.67449	1.07844	1.29333	1.42732	1.53025	1.61142	1.67812	1.73456	1.78335	1.86446	1.93013	1.98514	2.03235	2.07362			
	.90	1.28155	1.61749	1.79626	1.91655	2.00651	2.07785	2.13603	2.18354	2.23044	2.30307	2.35218	2.41190	2.45470	2.49224			
	.95	1.64465	1.94560	2.10727	2.21735	2.30056	2.36540	2.42058	2.46749	2.50793	2.57564	2.63092	2.67750	2.71770	2.75299			
	.99	2.32535	2.57215	2.70778	2.80077	2.87118	2.92765	2.97470	3.01455	3.05007	3.10312	3.15755	3.19852	3.23338	3.26521			

The entry in each case is the value of y^* satisfying $G_{k,v}(y^*, \rho) = p^* = 1 - \alpha$, where

$G_{k,v}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

ν	P^*	K	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
$\rho = .3$																
15.	.75		.69120	1.09341	1.30498	1.44625	1.55129	1.63440	1.70388	1.76093	1.81121	1.85494	1.95288	2.01987	2.06384	2.11159
	.90		1.34061	1.70091	1.83670	2.03620	2.13079	2.21116	2.27791	2.32483	2.36448	2.46767	2.53571	2.55315	2.64281	2.68648
	.95		1.75305	2.05588	2.25914	2.41703	2.51636	2.59613	2.66284	2.71956	2.76276	2.92120	2.97904	3.02302	3.07295	
	.99		2.60340	2.82704	3.11190	3.24032	3.33939	3.41757	3.48365	3.54083	3.59076	3.67524	3.74513	3.80480	3.85635	3.90336
16.	.75		.65013	1.09125	1.30210	1.44220	1.54740	1.63013	1.69829	1.75606	1.80609	1.85340	1.95635	2.01368	2.06239	2.10502
	.90		1.35375	1.69446	1.83502	2.03137	2.12104	2.20065	2.26673	2.32311	2.37219	2.45446	2.52172	2.57849	2.62752	2.67062
	.95		1.74559	2.05553	2.27344	2.40271	2.50074	2.57945	2.64504	2.70117	2.75016	2.82250	2.90000	2.95705	3.00633	3.04976
	.99		2.58344	2.90273	3.02400	3.21009	3.30642	3.38417	3.44923	3.50512	3.55406	3.63674	3.70495	3.76301	3.81357	3.85839
17.	.75		.63920	1.08932	1.29516	1.43978	1.54338	1.62638	1.69426	1.75179	1.80159	1.84453	1.95181	2.00824	2.05672	2.09914
	.90		1.33338	1.68506	1.82500	2.01364	2.11250	2.19144	2.25655	2.31282	2.35144	2.44593	2.50952	2.56569	2.61420	2.65683
	.95		1.73361	2.07616	2.26234	2.39020	2.48711	2.56488	2.62967	2.68511	2.73348	2.81479	2.88144	2.93778	2.98550	3.02936
	.99		2.56590	2.88123	3.05555	3.16353	3.27833	3.35481	3.41880	3.47375	3.52185	3.60304	3.66594	3.72678	3.77619	3.81989
18.	.75		.68236	1.08771	1.29732	1.43710	1.54095	1.62306	1.69059	1.74800	1.79761	1.85022	1.94723	2.00342	2.05169	2.09393
	.90		1.33039	1.68429	1.82634	2.00651	2.10497	2.18331	2.24931	2.30374	2.35196	2.43277	2.49877	2.55444	2.60249	2.64472
	.95		1.73407	2.06804	2.25256	2.37918	2.47510	2.55204	2.61614	2.67096	2.71850	2.79919	2.86507	2.92077	2.96893	3.01131
	.99		2.55226	2.83250	3.03799	3.16016	3.25349	3.32882	3.39185	3.44597	3.49334	3.57325	3.63905	3.69492	3.74342	3.78626
19.	.75		.68762	1.08622	1.29532	1.43471	1.53824	1.62010	1.68750	1.74462	1.79406	1.87637	1.94313	1.99911	2.04721	2.08928
	.90		1.32773	1.63004	1.87105	2.00073	2.09826	2.17603	2.24054	2.29566	2.34354	2.42374	2.48923	2.54445	2.59211	2.63398
	.95		1.72913	2.06083	2.24587	2.36939	2.46443	2.54065	2.60412	2.65841	2.70576	2.78533	2.85053	2.90565	2.95332	2.99526
	.99		2.53947	2.84552	3.01884	3.13934	3.23139	3.30568	3.36784	3.42121	3.46791	3.54670	3.61155	3.66659	3.71434	3.75649
20.	.75		.68555	1.08468	1.29352	1.43356	1.53582	1.61744	1.68464	1.74150	1.79087	1.87292	1.93946	1.99525	2.04318	2.08511
	.90		1.32534	1.67224	1.86630	1.99529	2.09226	2.16961	2.23376	2.28844	2.33601	2.41566	2.48070	2.53552	2.58284	2.62439
	.95		1.72472	2.05428	2.23610	2.36664	2.45490	2.53047	2.59339	2.64719	2.69414	2.77255	2.83754	2.89214	2.93935	2.98099
	.99		2.52797	2.83054	3.00174	3.12072	3.21160	3.28495	3.34632	3.39901	3.44512	3.52290	3.58690	3.64120	3.68831	3.72987
24.	.75		.63465	1.08057	1.28766	1.42531	1.52819	1.60903	1.67565	1.73204	1.78084	1.86205	1.92789	1.98306	2.03048	2.07195
	.90		1.31784	1.65420	1.85143	1.97322	2.07344	2.14933	2.21223	2.26381	2.31240	2.39038	2.45450	2.50761	2.55385	2.59444
	.95		1.71083	2.03418	2.21180	2.33329	2.42513	2.49368	2.55987	2.61216	2.65975	2.73429	2.78657	2.84593	2.89571	2.93598
	.99		2.45216	2.78091	2.94844	3.06622	3.14977	3.22008	3.27890	3.32939	3.37358	3.44811	3.50943	3.56144	3.60655	3.64633

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*, \rho) = P^* = 1 - \alpha$, where

$G_{k,\nu}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

 $\rho = .3$ (cont.)

ν	P^*	K	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
30.	.75	.63276	1.07542	1.27225	1.41512	1.52053	1.60079	1.66673	1.72252	1.77089	1.85127	1.91641	1.97100	2.01788	2.05888	
	.50	1.21042	1.63551	1.80236	1.91145	2.01489	2.12335	2.21012	2.24533	2.29516	2.36550	2.42775	2.43017	2.52536	2.56503	
	.55	1.65728	2.01453	2.16757	2.30642	2.35334	2.46753	2.52703	2.57785	2.62214	2.69644	2.75725	2.80859	2.85296	2.89198	
	.59	2.45726	2.73653	2.85853	3.00566	3.08938	3.15664	3.21287	3.26113	3.30335	3.37454	3.43311	3.48278	3.52585	3.56384	
36.	.75	.68137	1.07371	1.27354	1.41459	1.51562	1.59530	1.66084	1.71632	1.76431	1.84413	1.90881	1.96301	2.00354	2.05023	
	.50	1.20551	1.64474	1.82713	1.95032	2.04268	2.11620	2.17706	2.22936	2.27366	2.34912	2.41047	2.46211	2.50662	2.54568	
	.55	1.68230	2.00133	2.17422	2.28639	2.37681	2.44710	2.50551	2.55537	2.59379	2.67164	2.73121	2.78150	2.82495	2.86314	
	.59	2.43449	2.70853	2.88272	2.95305	3.05002	3.11526	3.16978	3.21654	3.25744	3.32639	3.38309	3.43117	3.47285	3.50961	
48.	.75	.67364	1.07027	1.27393	1.40920	1.50941	1.58349	1.65351	1.70855	1.75614	1.83527	1.89938	1.95308	1.99918	2.03948	
	.50	1.25944	1.65512	1.81515	1.93662	2.02753	2.09993	2.15979	2.21072	2.25495	2.32887	2.38910	2.43978	2.48344	2.52175	
	.55	1.67722	1.95227	2.13063	2.25722	2.35323	2.42194	2.47699	2.52767	2.57005	2.64108	2.69915	2.74814	2.79044	2.82762	
	.59	2.40553	2.67260	2.82137	2.92350	3.00183	3.06463	3.11704	3.16196	3.20123	3.26736	3.32176	3.36784	3.40777	3.44298	
60.	.75	.67365	1.06521	1.27118	1.40532	1.50571	1.58443	1.64915	1.70391	1.75127	1.82999	1.89375	1.94716	1.99299	2.03305	
	.50	1.25522	1.65540	1.80805	1.92848	2.01651	2.08027	2.14954	2.19954	2.25371	2.31684	2.37640	2.42650	2.46967	2.50752	
	.55	1.67065	1.97574	2.14161	2.25437	2.33326	2.40704	2.46329	2.51125	2.55302	2.62299	2.68016	2.72838	2.77000	2.80657	
	.59	2.35012	2.65150	2.79704	2.89735	2.97556	3.03436	3.08601	3.12965	3.16815	3.23266	3.28565	3.33054	3.36943	3.40372	
120.	.75	.67654	1.06412	1.25571	1.35340	1.43335	1.57637	1.64048	1.69471	1.74159	1.81950	1.88257	1.93538	1.98070	2.02031	
	.50	1.26055	1.61603	1.79599	1.91237	2.00086	2.07115	2.12924	2.17861	2.21446	2.29302	2.35125	2.40022	2.44238	2.47934	
	.55	1.65763	1.95174	2.11504	2.22603	2.31172	2.37765	2.43232	2.47852	2.51574	2.58731	2.64270	2.69540	2.72938	2.76506	
	.59	2.35762	2.65070	2.74543	2.84541	2.91820	2.97665	3.02537	3.06705	3.10551	3.16477	3.21505	3.25761	3.29445	3.32690	
∞	.75	.67449	1.06007	1.25026	1.35265	1.43106	1.56333	1.63163	1.68553	1.73155	1.80909	1.87148	1.92371	1.96851	2.00765	
	.50	1.26155	1.60631	1.78012	1.88649	1.93336	2.05230	2.10923	2.15753	2.19950	2.25953	2.32645	2.37429	2.41546	2.45153	
	.55	1.64435	1.95047	2.05353	2.20420	2.23470	2.34932	2.40194	2.44718	2.48550	2.55223	2.60594	2.65113	2.69009	2.72429	
	.59	2.35535	2.65324	2.70521	2.75501	2.83449	2.92020	2.96656	3.00624	3.04034	3.09557	3.14662	3.18691	3.22176	3.25243	

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*, \rho) = P^* = 1 - \alpha$, where

$G_{k,\nu}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

 $\rho = .4$

ν	P^*	K	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
15.	.75		.63120	1.07073	1.23973	1.46227	1.50053	1.57334	1.64227	1.69542	1.74323	1.82123	1.89440	1.93735	1.98231	2.02257
	.90		1.34061	1.62433	1.82140	1.95704	2.09300	2.16377	2.23157	2.29506	2.35157	2.40942	2.47233	2.52645	2.57262	2.61315
	.95		1.73305	2.08310	2.33335	2.5015	2.68335	2.83017	2.95017	3.05343	3.14274	3.21874	3.28243	3.33534	3.38124	3.42056
	.99		2.46240	2.91833	3.09753	3.22133	3.31532	3.39127	3.45447	3.50659	3.55585	3.60337	3.70064	3.75591	3.80382	3.84607
16.	.75		.63013	1.06337	1.23702	1.45907	1.49704	1.57442	1.63908	1.69199	1.73393	1.81622	1.87910	1.93179	1.97703	2.01659
	.90		1.33579	1.67335	1.85403	1.99155	2.08390	2.15330	2.22112	2.27411	2.32019	2.36725	2.41512	2.45307	2.49184	2.53032
	.95		1.74535	2.09236	2.34303	2.50933	2.65542	2.78223	2.89039	2.98044	3.05343	3.11874	3.17590	3.22443	3.26469	3.30653
	.99		2.50344	2.95445	3.13701	3.26177	3.33438	3.39890	3.45109	3.49434	3.52935	3.55593	3.63323	3.71761	3.76465	3.80609
17.	.75		.62820	1.05213	1.22463	1.44625	1.48333	1.57093	1.63440	1.68610	1.73455	1.81182	1.87444	1.92690	1.97194	2.01133
	.90		1.33338	1.67333	1.85577	1.99320	2.07574	2.15025	2.21197	2.26453	2.31020	2.35651	2.40391	2.45138	2.49662	2.53932
	.95		1.73561	2.08337	2.33231	2.50457	2.65659	2.80411	2.93180	3.04423	3.14233	3.22592	3.29559	3.36229	3.42806	3.49289
	.99		2.56690	2.97323	3.14603	3.26583	3.32701	3.33040	3.39166	3.44413	3.48956	3.52706	3.63032	3.68584	3.73016	3.77096
18.	.75		.62636	1.05536	1.22552	1.44776	1.48539	1.56793	1.63113	1.68465	1.73094	1.80793	1.87031	1.92257	1.96744	2.00557
	.90		1.33059	1.66855	1.85187	1.97570	2.06862	2.14262	2.20390	2.25607	2.30140	2.34722	2.39304	2.43908	2.48594	2.53531
	.95		1.73407	2.08552	2.33284	2.50377	2.64510	2.81819	2.97655	3.03084	3.07905	3.12190	3.16394	3.20632	3.24917	3.29135
	.99		2.55256	2.95452	3.12489	3.24291	3.32280	3.35015	3.35556	3.41731	3.46252	3.50859	3.60100	3.65391	3.69952	3.73976
19.	.75		.62462	1.05353	1.22504	1.44754	1.48539	1.56521	1.62823	1.68157	1.72772	1.80445	1.86662	1.91871	1.96342	2.00252
	.90		1.32773	1.65432	1.84679	1.96932	2.06228	2.13532	2.19571	2.24854	2.29557	2.34338	2.39086	2.43813	2.48526	2.53554
	.95		1.72913	2.04335	2.22442	2.34436	2.43491	2.50735	2.56756	2.61956	2.66374	2.73886	2.80029	2.85215	2.89534	2.93632
	.99		2.53947	2.93792	3.10603	3.22354	3.21125	3.28266	3.34226	3.39337	3.43801	3.51313	3.57477	3.62693	3.67208	3.71183
20.	.75		.62295	1.05255	1.22365	1.44654	1.48435	1.56421	1.62723	1.68057	1.72482	1.80134	1.86332	1.91524	1.95982	1.99879
	.90		1.32534	1.64112	1.83423	1.95674	2.05061	2.12974	2.19028	2.24181	2.29357	2.34541	2.42240	2.47374	2.51798	2.55630
	.95		1.72472	2.04233	2.21638	2.33594	2.42579	2.49766	2.53737	2.60835	2.65275	2.72722	2.78311	2.83950	2.88339	2.92290
	.99		2.52737	2.92311	3.09326	3.21033	3.19196	3.26251	3.32142	3.37190	3.41601	3.49025	3.55118	3.60275	3.64739	3.68569
24.	.75		.62035	1.05031	1.22162	1.44468	1.48231	1.56223	1.62525	1.67861	1.72242	1.80134	1.86332	1.91524	1.95982	1.99879
	.90		1.31784	1.64595	1.82756	1.94851	2.03082	2.11067	2.17012	2.22669	2.28461	2.33991	2.39780	2.44811	2.49144	2.52945
	.95		1.71088	2.02235	2.19221	2.30955	2.39732	2.46730	2.53556	2.57521	2.61843	2.69029	2.75008	2.80003	2.84315	2.88104
	.99		2.45216	2.77702	2.93692	3.04751	3.13170	3.19946	3.25604	3.30453	3.34691	3.41825	3.47662	3.52642	3.56936	3.60718

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*,\rho) = P^* = 1-\alpha$, where

$G_{k,\nu}(y^*,\rho)$ is given by (2.3).

Table 4 (cont.)

 $\rho = .4$ (cont.)

ν	P^*	K	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
30.	.75	.62278	1.05449	1.21234	1.37733	1.47233	1.54743	1.60926	1.65152	1.70671	1.76180	1.84251	1.89352	1.93721	1.97540	
	.90	1.31042	1.63202	1.81359	1.92259	2.02170	2.09189	2.15026	2.19990	2.24259	2.31498	2.37359	2.42238	2.46533	2.50355	
	.95	1.63758	2.00332	2.17283	2.33255	2.50340	2.43771	2.49440	2.54275	2.58482	2.65330	2.71287	2.76142	2.80331	2.84011	
	.99	2.45723	2.73215	2.88514	2.99212	3.07287	3.113782	3.19204	3.23849	3.27508	3.34740	3.40350	3.45059	3.49212	3.52835	
35.	.75	.63137	1.05124	1.24485	1.37211	1.46771	1.54246	1.60368	1.65523	1.70074	1.77537	1.83579	1.88637	1.92977	1.96770	
	.90	1.30551	1.63049	1.81590	1.92155	2.00375	2.07931	2.13718	2.18621	2.23115	2.29581	2.35764	2.40927	2.44814	2.48484	
	.95	1.63220	1.99092	2.15503	2.31503	2.45108	2.41825	2.47397	2.52146	2.56278	2.63199	2.68849	2.73611	2.77720	2.81329	
	.99	2.45345	2.70252	2.89275	2.99303	3.03452	3.09762	3.15027	3.19937	3.23476	3.30105	3.35947	3.40153	3.44142	3.47655	
40.	.75	.67234	1.04350	1.24050	1.36729	1.46193	1.53521	1.59719	1.64877	1.69334	1.76739	1.82732	1.87749	1.92053	1.95813	
	.90	1.29544	1.62109	1.79312	1.90322	1.95513	2.06422	2.12101	2.16933	2.21115	2.28106	2.33793	2.38573	2.42688	2.46294	
	.95	1.67722	1.97721	2.15328	2.34677	2.32852	2.39423	2.44879	2.45524	2.53564	2.60327	2.65845	2.70455	2.74505	2.78026	
	.99	2.40655	2.66714	2.81210	2.91163	2.96761	3.04844	3.09917	3.14255	3.18050	3.24428	3.29560	3.34087	3.37920	3.41294	
60.	.75	.67660	1.04636	1.23791	1.35474	1.43856	1.53249	1.59321	1.64455	1.68993	1.76263	1.82228	1.87220	1.91501	1.95243	
	.90	1.29522	1.61553	1.76530	1.90118	1.98700	2.05513	2.11141	2.15922	2.20055	2.25582	2.32222	2.37553	2.41425	2.44993	
	.95	1.67062	1.98547	2.15530	2.33371	2.31515	2.38007	2.43368	2.47971	2.51556	2.58625	2.64066	2.68649	2.72601	2.76069	
	.99	2.39012	2.64607	2.78819	2.88531	2.95001	3.01951	3.06910	3.11153	3.14857	3.21086	3.26194	3.30515	3.34255	3.37547	
120.	.75	.67654	1.04234	1.23278	1.35688	1.45176	1.52509	1.58529	1.63819	1.68016	1.75318	1.81225	1.86160	1.90407	1.94110	
	.90	1.29523	1.60450	1.76260	1.90594	1.97022	2.03714	2.09240	2.13533	2.18000	2.24787	2.30304	2.34936	2.38925	2.42417	
	.95	1.65765	1.94764	2.10244	2.21933	2.28278	2.35207	2.40447	2.44308	2.48786	2.55271	2.60558	2.65008	2.68845	2.72211	
	.99	2.35732	2.60451	2.74132	2.81539	2.90603	2.93232	3.01029	3.05080	3.08614	3.14551	3.19416	3.23525	3.27086	3.30216	
∞	.75	.67449	1.03874	1.22766	1.35237	1.44503	1.51776	1.57745	1.62730	1.67148	1.74332	1.80232	1.85126	1.89322	1.92987	
	.90	1.23155	1.55122	1.76543	1.87073	1.93367	2.01942	2.07356	2.11970	2.15590	2.26513	2.28016	2.32556	2.36459	2.39376	
	.95	1.64485	1.93593	2.08197	2.18540	2.26231	2.32458	2.37562	2.41504	2.45675	2.51979	2.57114	2.61435	2.65158	2.68422	
	.99	2.32655	2.56462	2.65593	2.73575	2.83361	2.90801	2.95323	2.99182	3.02556	3.08211	3.12840	3.16750	3.20130	3.23103	

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*, \rho) = P^* = 1 - \alpha$, where

$G_{k,\nu}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

ν	P^*	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
$\rho = .5$															
15.	.75	.69120	1.04441	1.22323	1.37143	1.44218	1.51373	1.57259	1.62236	1.66539	1.73691	1.79482	1.84330	1.88491	1.92127
	.90	1.34061	1.63548	1.84048	1.97957	2.04719	2.11757	2.17579	2.22529	2.25827	2.34008	2.39255	2.44773	2.49009	2.52723
	.95	1.75705	2.05791	2.25102	2.35623	2.44757	2.51527	2.57382	2.62387	2.66738	2.74028	2.79584	2.85008	2.89340	2.93148
	.99	2.60240	2.90713	3.07260	3.15577	3.22897	3.30663	3.41629	3.46722	3.51161	3.59607	3.64656	3.69835	3.74272	3.78171
16.	.75	.63013	1.04245	1.22333	1.34846	1.43598	1.51018	1.56878	1.61835	1.66120	1.72242	1.79007	1.83834	1.87975	1.91595
	.90	1.35773	1.63317	1.83244	1.95974	2.03335	2.10335	2.16607	2.21515	2.25775	2.32851	2.36684	2.43556	2.47751	2.51429
	.95	1.75525	2.05313	2.22718	2.34214	2.43049	2.50025	2.55814	2.60750	2.65046	2.72243	2.78120	2.83075	2.87351	2.91106
	.99	2.56244	2.83312	3.05110	3.17701	3.25377	3.32523	3.38358	3.43415	3.47789	3.55130	3.61135	3.66204	3.70582	3.74429
17.	.75	.60220	1.04072	1.22412	1.34575	1.43508	1.50704	1.56544	1.61483	1.65752	1.72847	1.79590	1.83058	1.87523	1.91128
	.90	1.35120	1.63553	1.83727	1.95772	2.03033	2.10027	2.15757	2.20627	2.24854	2.31914	2.37660	2.42931	2.46651	2.50298
	.95	1.75161	2.04918	2.21622	2.33104	2.41810	2.48713	2.54440	2.59322	2.63570	2.70625	2.76494	2.81391	2.85616	2.89327
	.99	2.56330	2.83215	3.02753	3.14161	3.22816	3.29763	3.35549	3.40496	3.44810	3.52052	3.57980	3.62984	3.67307	3.71106
18.	.75	.60238	1.03319	1.22215	1.34355	1.43342	1.50426	1.56247	1.61171	1.65426	1.72497	1.78220	1.83012	1.87122	1.90715
	.90	1.35039	1.63003	1.82182	1.94764	2.02423	2.09314	2.15005	2.19843	2.24041	2.31051	2.36756	2.41552	2.45581	2.49300
	.95	1.75407	2.04040	2.20762	2.32151	2.40719	2.47357	2.53229	2.58064	2.62271	2.69314	2.75064	2.79910	2.84091	2.87761
	.99	2.55236	2.84533	3.00681	3.11927	3.20453	3.27315	3.33024	3.37906	3.42164	3.49314	3.55168	3.60112	3.64384	3.68139
19.	.75	.60782	1.03763	1.22038	1.34150	1.43113	1.50178	1.55983	1.60933	1.65126	1.72186	1.77891	1.82667	1.86765	1.90346
	.90	1.32773	1.64367	1.81637	1.93217	2.01834	2.08679	2.14337	2.19145	2.23318	2.30284	2.35952	2.40717	2.44818	2.48413
	.95	1.72513	2.03250	2.19550	2.31251	2.39743	2.46531	2.52155	2.56948	2.61118	2.68090	2.73795	2.78556	2.82738	2.86374
	.99	2.55947	2.82753	2.95853	3.05542	3.13370	3.20136	3.26773	3.32535	3.37801	3.44686	3.50389	3.55223	3.59402	3.63076
20.	.75	.60555	1.02651	1.21880	1.33955	1.42907	1.49936	1.55746	1.60543	1.64875	1.71906	1.77596	1.82359	1.86444	1.90015
	.90	1.32534	1.64248	1.81262	1.92727	2.01301	2.08111	2.13738	2.18521	2.22570	2.29597	2.35233	2.39969	2.44046	2.47620
	.95	1.72472	2.02733	2.19250	2.30446	2.39233	2.45614	2.51195	2.55951	2.60087	2.67012	2.72662	2.77423	2.81530	2.85135
	.99	2.53797	2.81273	2.97155	3.06169	3.15498	3.23185	3.28757	3.33524	3.37682	3.44658	3.50369	3.55223	3.59402	3.63076
24.	.75	.60485	1.02375	1.21382	1.33384	1.42761	1.49256	1.55000	1.59957	1.64054	1.71026	1.76666	1.81388	1.85435	1.88973
	.90	1.31734	1.63160	1.79358	1.91191	1.99229	2.06328	2.11861	2.16562	2.20640	2.27444	2.32977	2.37627	2.41623	2.45134
	.95	1.71628	2.00001	2.16735	2.27350	2.33175	2.38178	2.42718	2.46156	2.49536	2.56320	2.63126	2.73764	2.77762	2.81271
	.99	2.48216	2.75727	2.92072	3.02657	3.10552	3.17026	3.22447	3.27034	3.31035	3.37758	3.43267	3.47923	3.51949	3.55490

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*, \rho) = p^* = 1 - \alpha$, where $G_{k,\nu}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)
 $\rho = .5$ (cont.)

ν	P^*	K	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
30.	.75		.62276	1.02332	1.20327	1.32607	1.41320	1.46332	1.54261	1.59079	1.63341	1.70154	1.75745	1.80423	1.84435	1.87940
	.90		1.31042	1.62007	1.77552	1.89876	1.97582	2.04571	2.10012	2.14633	2.18539	2.25323	2.30756	2.35320	2.39246	2.42686
	.95		1.63726	1.92903	2.14730	2.29461	2.39333	2.45256	2.49722	2.53718	2.60299	2.65664	2.70181	2.74075	2.77491	
	.99		2.45726	2.72553	2.87084	2.97247	3.04950	3.11131	3.16279	3.20622	3.24523	3.30976	3.36262	3.40730	3.44594	3.47993
36.	.75		.68137	1.02528	1.20550	1.32426	1.41196	1.46103	1.53772	1.58564	1.62304	1.69577	1.75136	1.79766	1.83774	1.87258
	.90		1.39551	1.61274	1.77266	1.88678	1.96826	2.03414	2.08794	2.13362	2.17322	2.23927	2.29294	2.33802	2.37679	2.41074
	.95		1.63330	1.97656	2.13237	2.23341	2.31776	2.38095	2.43327	2.47781	2.51651	2.58121	2.63395	2.67833	2.71658	2.75013
	.99		2.45245	2.69421	2.83355	2.93735	3.01235	3.07249	3.12256	3.16538	3.20372	3.26545	3.31682	3.36023	3.39777	3.43078
48.	.75		.67564	1.02322	1.20153	1.31952	1.40670	1.47533	1.53165	1.57925	1.62036	1.68861	1.74320	1.78996	1.82953	1.86410
	.90		1.23522	1.46363	1.60555	1.67444	1.73554	2.01934	2.07389	2.11791	2.15694	2.22200	2.27436	2.31924	2.35740	2.39082
	.95		1.67122	1.98117	2.14822	2.21624	2.28228	2.35324	2.40951	2.45314	2.49104	2.55439	2.60599	2.64941	2.68682	2.71962
	.99		2.40652	2.65632	2.75860	2.83439	2.89633	3.02499	3.07334	3.11467	3.15069	3.21119	3.26072	3.30256	3.33872	3.37052
60.	.75		.67680	1.02134	1.19910	1.31670	1.40356	1.47193	1.52803	1.57544	1.61638	1.68434	1.73929	1.78525	1.82464	1.85905
	.90		1.23522	1.46363	1.60555	1.67444	1.73554	2.01934	2.07389	2.11791	2.15694	2.22200	2.27436	2.31924	2.35740	2.39082
	.95		1.67065	1.98205	2.14822	2.21624	2.28228	2.35324	2.40951	2.45314	2.49104	2.55439	2.60599	2.64941	2.68682	2.71962
	.99		2.55012	2.65313	2.77319	2.86911	2.94014	2.99704	3.04438	3.08482	3.12508	3.17926	3.22769	3.26860	3.30395	3.33504
120.	.75		.67654	1.01733	1.19423	1.31103	1.37732	1.43513	1.52065	1.56783	1.61643	1.67537	1.73024	1.77569	1.81493	1.84902
	.90		1.23623	1.46355	1.60555	1.67444	1.73554	2.01934	2.07389	2.11791	2.15694	2.22200	2.27436	2.31924	2.35740	2.39082
	.95		1.67755	1.93403	2.08201	2.14822	2.21624	2.28228	2.35324	2.40951	2.45314	2.49104	2.55439	2.60599	2.64941	2.68682
	.99		2.55722	2.59742	2.72223	2.81583	2.87780	2.94233	2.98772	3.02545	3.06519	3.11680	3.16310	3.20218	3.23594	3.26562
∞	.75		.67449	1.01387	1.18943	1.30752	1.37114	1.43349	1.51373	1.56038	1.60065	1.66748	1.72148	1.76632	1.80531	1.83909
	.90		1.23155	1.46355	1.60555	1.67444	1.73554	2.01934	2.07389	2.11791	2.15694	2.22200	2.27436	2.31924	2.35740	2.39082
	.95		1.64485	1.91633	2.06208	2.16033	2.23382	2.29219	2.34044	2.38144	2.41702	2.47611	2.52473	2.56534	2.60029	2.63093
	.99		2.58635	2.55722	2.68465	2.77156	2.83598	2.89327	2.93271	2.96978	3.00306	3.03616	3.10040	3.13772	3.16993	3.19823

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*, \rho) = p^* = 1 - \alpha$, where

$G_{k,\nu}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

ν	P^*	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
15.	.75	1.6120	1.0172	1.1003	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.50	1.5669	1.0410	1.1000	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.25	1.7220	2.0000	2.1000	2.2015	2.3490	2.4553	2.5152	2.5713	2.6334	2.6910	2.7140	2.8240	2.8720	2.90479
	.00	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000
16.	.75	1.6120	1.0172	1.1003	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.50	1.5669	1.0410	1.1000	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.25	1.7220	2.0000	2.1000	2.2015	2.3490	2.4553	2.5152	2.5713	2.6334	2.6910	2.7140	2.8240	2.8720	2.90479
	.00	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000
17.	.75	1.6120	1.0172	1.1003	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.50	1.5669	1.0410	1.1000	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.25	1.7220	2.0000	2.1000	2.2015	2.3490	2.4553	2.5152	2.5713	2.6334	2.6910	2.7140	2.8240	2.8720	2.90479
	.00	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000
18.	.75	1.6120	1.0172	1.1003	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.50	1.5669	1.0410	1.1000	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.25	1.7220	2.0000	2.1000	2.2015	2.3490	2.4553	2.5152	2.5713	2.6334	2.6910	2.7140	2.8240	2.8720	2.90479
	.00	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000
19.	.75	1.6120	1.0172	1.1003	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.50	1.5669	1.0410	1.1000	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.25	1.7220	2.0000	2.1000	2.2015	2.3490	2.4553	2.5152	2.5713	2.6334	2.6910	2.7140	2.8240	2.8720	2.90479
	.00	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000
20.	.75	1.6120	1.0172	1.1003	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.50	1.5669	1.0410	1.1000	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.25	1.7220	2.0000	2.1000	2.2015	2.3490	2.4553	2.5152	2.5713	2.6334	2.6910	2.7140	2.8240	2.8720	2.90479
	.00	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000
24.	.75	1.6120	1.0172	1.1003	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.50	1.5669	1.0410	1.1000	1.2015	1.3490	1.4553	1.4910	1.5152	1.5713	1.6334	1.6140	1.7240	1.7720	1.80479
	.25	1.7220	2.0000	2.1000	2.2015	2.3490	2.4553	2.5152	2.5713	2.6334	2.6910	2.7140	2.8240	2.8720	2.90479
	.00	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000

The entry in each case is the value of y^* satisfying $G_{k,v}(y^*,) = P^* = 1 - \alpha$, where

$G_{k,v}(y^*,)$ is given by (2.3).

Table 4 (cont.)

 $\rho = .6$ (cont.)

ν	$P^* K$	1.	2.	3.	4.	5.	6.	7.	8.	9.	11.	13.	15.	17.	19.
30.	.75	.62276	.53373	1.11324	1.21973	1.27038	1.41317	1.45783	1.52215	1.59570	1.67991	1.69883	1.70048	1.73555	1.76906
	.50	1.11042	1.53737	1.71120	1.71220	1.67220	1.52310	2.03255	2.03943	2.11533	2.17760	2.21593	2.23255	2.30391	2.33503
	.25	1.53728	1.53041	2.11151	2.21038	2.21031	2.35220	2.35910	2.44081	2.47865	2.51277	2.53571	2.53571	2.63216	2.69330
	.05	2.15722	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037	2.21037
36.	.75	.62157	.53253	1.11200	1.21848	1.26913	1.41192	1.45658	1.52090	1.59445	1.67866	1.69758	1.70023	1.73530	1.76881
	.50	1.11011	1.53706	1.71089	1.71189	1.67189	1.52279	2.03224	2.03912	2.11502	2.17729	2.21562	2.23224	2.30360	2.33472
	.25	1.53705	1.53018	2.11128	2.21015	2.21008	2.35203	2.35893	2.44064	2.47848	2.51260	2.53554	2.53554	2.63199	2.69313
	.05	2.15715	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030	2.21030
48.	.75	.62134	.53230	1.11177	1.21825	1.26890	1.41169	1.45634	1.52066	1.59421	1.67842	1.69734	1.70000	1.73507	1.76858
	.50	1.11000	1.53695	1.71078	1.71178	1.67178	1.52268	2.03213	2.03901	2.11491	2.17718	2.21551	2.23213	2.30349	2.33461
	.25	1.53694	1.53007	2.11117	2.21004	2.20997	2.35192	2.35882	2.44053	2.47837	2.51249	2.53543	2.53543	2.63188	2.69302
	.05	2.15705	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020	2.21020
60.	.75	.62111	.53207	1.11154	1.21802	1.26867	1.41146	1.45611	1.52043	1.59398	1.67819	1.69711	1.70026	1.73533	1.76884
	.50	1.10987	1.53682	1.71065	1.71165	1.67165	1.52255	2.03200	2.03888	2.11478	2.17705	2.21540	2.23202	2.30338	2.33450
	.25	1.53681	1.52994	2.11104	2.20991	2.20984	2.35185	2.35875	2.44046	2.47830	2.51242	2.53536	2.53536	2.63181	2.69295
	.05	2.15695	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010	2.21010
120.	.75	.62088	.53184	1.11131	1.21779	1.26844	1.41113	1.45578	1.52010	1.59365	1.67786	1.69678	1.70023	1.73530	1.76881
	.50	1.10964	1.53659	1.71052	1.71152	1.67152	1.52242	2.03187	2.03875	2.11465	2.17692	2.21530	2.23192	2.30328	2.33440
	.25	1.53658	1.52971	2.11089	2.20976	2.20969	2.35176	2.35866	2.44037	2.47821	2.51233	2.53527	2.53527	2.63172	2.69286
	.05	2.15672	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005	2.21005

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*, \rho) = P^* = 1 - \alpha$, where

$G_{k,\nu}(y^*, \rho)$ is given by (2.3).

Table 4 (cont.)

 $\rho = .3$

ν	p^*	1.	2.	3.	4.	5.	5.	6.	7.	8.	9.	11.	13.	15.
15.	.50	1.51091	1.51097	1.51174	1.51293	1.51431	1.51593	1.51700	1.51700	1.51700	1.51724	2.02473	2.02253	2.09449
15.	.55	1.75535	1.75570	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
17.	.50	1.52713	1.52713	1.52713	1.52713	1.52713	1.52713	1.52713	1.52713	1.52713	1.52713	2.02473	2.02253	2.09449
17.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
20.	.50	1.55113	1.55113	1.55113	1.55113	1.55113	1.55113	1.55113	1.55113	1.55113	1.55113	2.02473	2.02253	2.09449
20.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
25.	.50	1.57223	1.57223	1.57223	1.57223	1.57223	1.57223	1.57223	1.57223	1.57223	1.57223	2.02473	2.02253	2.09449
25.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
30.	.50	1.59113	1.59113	1.59113	1.59113	1.59113	1.59113	1.59113	1.59113	1.59113	1.59113	2.02473	2.02253	2.09449
30.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
35.	.50	1.60713	1.60713	1.60713	1.60713	1.60713	1.60713	1.60713	1.60713	1.60713	1.60713	2.02473	2.02253	2.09449
35.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
40.	.50	1.62113	1.62113	1.62113	1.62113	1.62113	1.62113	1.62113	1.62113	1.62113	1.62113	2.02473	2.02253	2.09449
40.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
45.	.50	1.63513	1.63513	1.63513	1.63513	1.63513	1.63513	1.63513	1.63513	1.63513	1.63513	2.02473	2.02253	2.09449
45.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431
50.	.50	1.64913	1.64913	1.64913	1.64913	1.64913	1.64913	1.64913	1.64913	1.64913	1.64913	2.02473	2.02253	2.09449
50.	.55	1.77011	1.77011	2.11071	2.11270	2.27431	2.28333	2.33225	2.33225	2.33225	2.33450	2.44253	2.44155	2.51431

The entry in each case is the value of y^* satisfying $G_{k,\nu}(y^*, r) = p^* = 1 - \alpha$,

where $G_{k,\nu}(y^*, r)$ is given by (2.3).

6. APPLICATIONS

In this section we discuss briefly some specific applications illustrating the use of the tables. These applications relate to selection procedures, multiple comparisons, prediction intervals and tests of hypotheses.

6.1 Selection procedures

Let π_1, \dots, π_k be k independent normal populations with unknown means μ_1, \dots, μ_k , respectively, and common unknown variance σ^2 . Gupta [5], [7] proposed a procedure R_1 for selecting a nonempty subset from the k populations so that the population associated with the largest μ_i is included in the selected subset (this event is called a correct selection (CS)) with a probability at least equal to P^* ($1/k < P^* < 1$). Let $Y_i = \bar{X}_i$, $i = 1, \dots, k$, be the means of samples of size n from these populations. The rule R_1 is: Select π_i if and only if $Y_i \geq \max_{1 \leq j \leq k} Y_j - \frac{ds}{\sqrt{n}}$, where s^2 is the usual pooled estimator of σ^2 based on all the k samples such that vs^2/σ^2 has a χ^2 distribution with $v = k(n-1)$ degrees of freedom. The infimum of PCS, the probability of a CS, occurs when $\mu_1 = \dots = \mu_k$ for all values of $\sigma^2 > 0$: Thus the value of d satisfying the probability requirement is given by $G_{k-1, v}(d/\sqrt{2}; 1/2) = P^*$. Therefore, $d = \sqrt{2} y^*(1-P^*, k-1, k(n-1), 0.5)$.

Suppose the above k populations represent experimental categories and they are compared with a control population π_0 which is $N(\mu_0, \sigma^2)$ where μ_0 is unknown. Let $Y_0 = \bar{X}_0$ be the mean of a sample of size m from π_0 . The pooled estimator s^2 is now based on $v = k(n-1) + (m-1)$ degrees of freedom. Gupta and Sobel [10] proposed a rule R_2 for selecting a subset of the k experimental populations such that all populations better than the control (i.e. those for which $\mu_i \geq \mu_0$) are included in the selected subset with a probability at least equal to P^* . We propose a modified procedure R_2' :

Select π_i if and only if $\bar{X}_i \geq \bar{X}_0 - Ds \sqrt{\frac{1}{m} + \frac{1}{n}}$, where $D > 0$ is to be determined to satisfy the probability requirement. It is easy to show that the infimum of PCS is equal to $G_{k,v}(D; \rho)$ where $\rho = n/(n+m)$. This gives $D = y^*(1-P^*, k, v, \rho)$.

6.2 Multiple comparisons between k treatment means and a control mean

The problem of the simultaneous one-sided comparisons of the means of k treatments with that of a control group has been discussed by Dunnett [3]. Let $\bar{X}_0, \bar{X}_1, \dots, \bar{X}_k$ be independent sample means based on m, n, \dots, n independent observations from normal populations with unknown means $\mu_0, \mu_1, \dots, \mu_k$, respectively, and a common unknown variance σ^2 . Let s^2 be the usual pooled unbiased estimator of σ^2 based on $v = k(n-1) + (m-1)$ degrees of freedom such that vs^2/σ^2 has a χ^2 -distribution with v degrees of freedom. Then a set of k exact one-sided (lower) 100% simultaneous confidence intervals for the differences $\mu_i - \mu_0$, $i = 1, \dots, k$, is given by

$$(\bar{X}_i - \bar{X}_0) - y^*(1-\gamma, k, v, \rho) s \sqrt{\frac{1}{m} + \frac{1}{n}}, \quad i = 1, \dots, k$$

where $\rho = n/(n+m)$.

6.3 Prediction intervals to contain all of k future means

Let Y_1, \dots, Y_n be the n observations of a current sample from a normal population with mean μ and variance σ^2 , both unknown. Let \bar{Y} be the sample mean and s^2 be the usual unbiased estimator of σ^2 on $v = n-1$ degrees of freedom. Let $\bar{X}_1, \dots, \bar{X}_k$ be the (unknown) sample means of k additional future independent samples each of size m. The \bar{X}_i are independent of \bar{Y} and s^2 . An exact lower simultaneous 100% prediction interval for the sample means $\bar{X}_1, \dots, \bar{X}_k$ is given by $\bar{Y} - y^*(1-\gamma, k, v, \rho) s \sqrt{\frac{1}{m} + \frac{1}{n}}$, where $\rho = m/(m+n)$.

6.4 Test procedures for multiple comparisons

Krishnaiah [12] proposed test procedures for the multiple comparisons of means (mean vectors) against one-sided alternatives under a general analysis of variance (multivariate analysis of variance) model. Let us consider k normal populations with unknown means μ_1, \dots, μ_k and a common unknown variance σ^2 . Consider the problem of testing

$H_i: c_i' \mu = 0$ simultaneously against the respective alternatives

$K_i: c_i' \mu > 0, i = 1, \dots, q$ ($q < k$), where $\mu' = (\mu_1, \dots, \mu_k)$,

and the c_i are $k \times 1$ vectors of known elements such that $c_i' c_i = 1$,

$c_i' \underline{1} = 0$ for $i = 1, \dots, q$, and $c_i' c_j = \rho$ (> 0) for $i \neq j = 1, \dots, q$. Here $\underline{1}$ denotes a column vector whose elements are unity. It is required that

$$\text{pr}(\text{any } H_i \text{ is rejected} \mid \bigcap_{i=1}^q H_i) = \alpha.$$

Let $\bar{Y}_i, i = 1, \dots, k$, the sample means based on n independent observations from each population and let s^2 denote the usual pooled unbiased estimator of σ^2 on $v = k(n-1)$ degrees of freedom. Then, as a special case of the procedure proposed by Krishnaiah [12], we use the procedure:

$$\text{Reject } H_i \text{ (against } K_i) \text{ if } t_i = \frac{\sqrt{n} c_i' \bar{Z}}{s} > t_{\alpha}$$

where $\bar{Z} = (\bar{Y}_1, \dots, \bar{Y}_k)$ and t_{α} satisfying the error restriction is given by $t_{\alpha} = y^*(1-\alpha, k, v, \rho)$.

REFERENCES

- [1] Bechhofer, R. E. (1954). A single-sample multiple decision procedure for ranking means of normal populations with known variances. Ann. Math. Statist., 25, 16-39.
- [2] Cornish, E. A. (1954). The multivariate t-distribution associated with a set of normal sample deviates. Austral. J. Phys., 7, 531-542.
- [3] Dunnett, C. W. (1955). A multiple comparison procedure for comparing several treatments with a control. J. Amer. Statist. Assoc., 50, 1096-1121.
- [4] Dunnett, C. W. and Sobel, M. (1954). A bivariate generalization of Student's t-distribution with tables for certain special cases. Biometrika, 41, 153-169.
- [5] Gupta, S. S. (1956). On a decision rule for a problem in ranking means. Ph.D. Thesis (Mimeo. Ser. No. 150), Inst. of Statistics, University of North Carolina, Chapel Hill.
- [6] Gupta, S. S. (1963). Probability integrals of multivariate normal and multivariate t. Ann. Math. Statist., 34, 792-828.
- [7] Gupta, S. S. (1965). On some multiple decision (selection and ranking) rules. Technometrics, 7, 225-245.
- [8] Gupta, S. S., Nagel, K. and Panchapakesan, S. (1973). On the order statistics from equally correlated normal random variables. Biometrika, 60, 403-413.
- [9] Gupta, S. S. and Sobel, M. (1957). On a statistic which arises in selection and ranking problems. Ann. Math. Statist., 28, 957-967.
- [10] Gupta, S. S. and Sobel, M. (1958). On selecting a subset that contains all populations better than a standard. Ann. Math. Statist., 29, 235-244.
- [11] Hartley, H. O. (1944). Studentization or the elimination of the standard deviation of the parent population from the random-sample distribution of statistics. Biometrika, 33, 173-180.
- [12] Krishnaiah, P. R. (1965). Multiple comparison tests in multi-response experiments. Sankhyā, Ser. A, 27, 65-72.
- [13] Krishnaiah, P. R. and Armitage, J. V. (1966). Tables for multivariate t distribution. Sankhyā, Ser. B, 28, 31-56.
- [14] Milton, R. C. (1963). Tables of equally correlated multivariate normal probability integral. Tech. Report No. 27, Dept. of Statistics, University of Minnesota, Minneapolis.
- [15] Pillai, K.C.S. and Ramachandran, K. V. (1954). On the distribution of the ratio of the i-th observation in an ordered sample from a normal population to an independent estimate of the standard deviation. Ann. Math. Statist., 25, 565-572.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report #83-31	2. GOVT ACCESSION NO. A131529	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) On the Distribution of the Studentized Maximum of Equally Correlated Normal Random Variables		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) Shanti S. Gupta, S. Panchapakesan, and Joong K. Sohn		6. PERFORMING ORG. REPORT NUMBER Technical Report #83-31
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics West Lafayette, IN 47907		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0455
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, DC		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 1983
		13. NUMBER OF PAGES 31
		15. SECURITY CLASS (of this report) Unclassified
		19a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Normal variables; Equicorrelated; Studentized; Maximum; multivariate t; Percentage points; Tables; Subset selection procedures; Multiple comparisons; Prediction intervals; Tests of hypotheses.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let X_1, \dots, X_k have a joint k-variate normal distribution with zero means, common unknown variance σ^2 and known correlation matrix (ρ_{ij}) , where $\rho_{ij} = \rho$ for all $i \neq j$. Let s^2 be distributed independently of the X_i such that vs^2/σ^2 has a chi-squared distribution with v degrees of freedom. New tables with wider coverage and more accuracy than the previously published ones are given for the percentage points of (over)		

DD FORM 1473
1 JAN 73

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

$Y = \max_{1 \leq i \leq k} \frac{X_i}{S}$. Some basic theoretical results are given in Section 2. The next

section describes Hartley's results for approximating the distribution function of Y . Besides a brief review of existing tables (Section 4), the paper discusses the construction of new tables based on Hartley's results (Section 5) and some specific applications (Section 6).

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

